


# Now There's Hope for Lasting Relief with Non-Surgical Spinal Decompression



## Are You Suffering From Chronic Back Pain?

Treat Back Pain and Sciatica caused by:

- ▶ Bulging/Herniated Discs
- ▶ Degenerative Disc Disease
- ▶ Facet Syndrome
- ▶ Failed Back Surgery

Without the use of:

- ✗ Drugs
- ✗ Injections
- ✗ Surgery

*"As a surgeon, I only want to do surgery when I absolutely have to. Non-Surgical Spinal Decompression Therapy gives my patients a more conservative treatment option that can eliminate the need for surgery altogether, and that's a very good thing."*

— Dr. Bernard Zeligler, DO, FRCGS, FRCR, FRC —  
Osteopathic Physician and Orthopedic Surgeon  
Founding Dean and Provost of Touro University  
College of Osteopathic Medicine, Vallejo, CA

All Pages In  
Printed Versions  
Are In Full Color

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient/educational purposes. Visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information.

## Inside This Special Report on Back Pain...



Learn about the many causes, forms, and symptoms of back pain, and how Non-Surgical Spinal Decompression can deliver real relief for back pain sufferers!



Are you considering surgery? Spinal decompression could be the non-invasive alternative you are looking for. Even post-surgical patients can benefit!



Are you tired of turning to medications that provide only temporary relief? See how you can find drug-free results with Non-Surgical Spinal Decompression.



Non-Surgical Spinal Decompression has been proven effective in several clinical studies. Thousands of patients a day are treated with this safe, non-surgical therapy.

This area may be customized with your clinic information. Otherwise it is left blank, and you may attach a business card, sticker, or use a stamp. See order form for customization details and pricing or call with inquiries at 877-968-8631.

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient/educational purposes. Visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information.

The testimonials given in this brochure are from actual patients who have undergone Non-Surgical Spinal Decompression Therapy. These testimonials have been provided by a consortium of practitioners utilizing Non-Surgical Spinal Decompression Therapy. Therefore, the testimonials appearing in this report may or may not be from the specific doctor(s)/office(s) providing this report. Testimonials may have been edited for length and clarity. To protect patient privacy, last names have been removed and modified. Photographs substituted.

Document No. 06N149P4-C-09-09

This report has not been evaluated by the FDA.

To order: [medwestpublications.com](http://medwestpublications.com), 1-877-968-8631

Copyright © 2005-2009 MediaWest. All Rights Reserved. [www.medwestpublications.com](http://www.medwestpublications.com)

# What Is Spinal Decompression Therapy?

## Proven Non-Surgical Treatment for Back Pain

**Non-Surgical Spinal Decompression (NSSD),** also referred to as Spinal Decompression, is proven to treat back pain and sciatica caused by bulging, herniated, and degenerative discs or facet syndrome. Even post-surgical patients and those suffering from certain types of stenosis (a narrowing of the spinal canal) have reported significant pain relief from NSSD treatments.

*Over a series of relaxing treatment sessions, patients can experience powerful pain reduction and remission of symptoms. Some patients even notice an improvement in their symptoms after the first few treatments!*

### Unlike Anything You've Ever Experienced

Non-Surgical Spinal Decompression, not to be confused with linear traction, gently lengthens and decompresses the spine, creating negative pressures<sup>1</sup> within the discs. This reversal of pressure creates an intradiscal vacuum that helps to reposition bulging discs and pull extruded disc material back into place.

1. Gustafson, MD, William Martin, MD, Effects of Vertebral Axial Decompression on Vertebral Fractures, *Journal of Neurology*, 81(3), 1904.  
2. Michael Schumala, MD, MD, PhD, Spinal Disc, *Neurology*, 46(4), 1000, 1996.



taking pressure off pinched nerves. Spinal experts believe that nutrients, oxygen, and fluids are drawn into the disc to create a revitalized environment conducive to healing.<sup>2</sup>

Experts surmise that Non-Surgical Spinal Decompression stimulates the body's repair mechanism, providing the building blocks needed to mend injured and degenerated discs. NSSD enables the body to get to work and heal itself! Now there truly is hope for lasting relief without drugs, injections, or surgery.

**"Driving, walking, sitting, standing, and bending were all impossible to do without unbearable pain. I met with the doctor offering Non-Surgical Spinal Decompression therapy and he felt very strongly that it could be of help. By this time I was skeptical of anything claiming its ability to relieve my pain. WOW! I can't believe the difference! I met many people at his office that had a poor outcome from their surgical procedures and am glad that I chose the route I did. I would highly recommend this therapy to anyone with a problem similar to mine." — Christopher R.**

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.urmcawebpublications.com](http://www.urmcawebpublications.com) or call 877-968-8631 for ordering information.

## Invention Born of Necessity

*How one physician's injury led to his invention of Non-Surgical Spinal Decompression.*

Allan Dyer, MD, PhD, is the inventor and founder of the revolutionary technology behind Non-Surgical Spinal Decompression. As former Deputy Minister of Health in Ontario, Canada, Dr. Dyer's many contributions to health sciences include his extensive, pioneering research that contributed to the development of the heart defibrillator.

Dr. Dyer's own experience with back pain began when he was debilitated by a herniated disc. After conventional treatments failed, his creative intelligence went into high gear. He developed a method of exerting pull on the spine that is far more sophisticated than traditional traction. He discovered that by slowly increasing pull-tension on the spine, followed by a hold then partial release, and repeating these steps over a 30–45 minute time period, a vacuum could be created within injured discs. This vacuum was found to assist in repositioning extruded disc material back into place, and has been credited with creating an environment within injured discs conducive to healing.

Dr. Dyer set about designing a device to implement this method. After more than six years in research and development with a team of physicians, engineers, and technicians at major teaching hospitals, Dr. Dyer introduced the WAX-D™ unit in 1991. WAX-D, short for Vertebral Axial Decompression, is the predicate device from which all Non-Surgical Spinal Decompression Therapy units are modeled.

Happily, Dr. Dyer used his invention on his own injured disc. Soon he was able to walk pain-free and has been doing so for more than 15 years. Today, thousands of Spinal Decompression units operate throughout the world, treating tens of thousands of patients a day.



### "I had pain in my back

and down my right leg for years. I was having trouble even walking up and down stairs. I had four epidurals, which didn't help much, and the last one was very painful. Lying on my stomach was almost impossible.

I started my mornings off with pain medication. The more I tried to do throughout the day, the worse I felt. For the past two years, I had to sleep in a recliner with a pillow underneath my legs, and even then it was hard to sleep.

I had seen two chiropractors, a physical therapist, and a pain management specialist, but nothing helped until I underwent Non-Surgical Spinal Decompression Therapy. Now I'm sleeping really well, and I never wake up due to back pain. I can finally do the things I want to do, without pain."

—Ruthann T.—

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information. Copyright © 2005–2009 MediaWest. All Rights Reserved.

## Four Years after Treatment: Patients Enjoy Long Term Results!

Pain Levels Based on Oswestry Pain Scale



Elapsed Time: 20 Treatment Sessions

A four-year, multi-center followup study\* on the lasting effects of Vertebral Axial Decompression (VAX-D) showed remarkable levels of sustained relief in 23 back pain patients. All had undergone several types of treatment before receiving VAX-D therapy. Before treatment, patients reported an average pain level of 7.41 out of 6 possible &

**Immediately following treatment protocol:**

- Average pain levels were reduced from 7.41 to 2.41 immediately after treatment.

- 71% showed more than a 50% reduction in pain.

**Four years following treatment protocol:**

- 91% were able to resume their normal daily activities.

- 86% showed a 50% or better pain reduction.

- Employment status increased by 40% among those previously out of work due to back pain.

- 87% were either working or were retired without having back pain as the cause for retirement.

- 52% of respondents reported a pain level of zero.

- Average pain levels were further reduced to 1.57.

**Pain relief not only lasted but improved!**

\* E. H. Odeh, MD, PhD and D. A. Sanderson, DO. In the March 2009 edition of *Spine* (Neurology News) (Vol. 38).

† This research was performed at a VAX-D unit, the International Institute for Non-Surgical Spinal Decompression, a non-profit research center for non-surgical spinal decompression treatment of NCD. © Medical Technologies.



**"I have given close to**

10,000 Non-Surgical Spinal Decompression treatments over the last three years. Knowing how many people this helps, I would never practice again without this technology."

— **Dr. Adrian Marcus, DC** —

Chiropractic Physician

**"My pain was so bad**

that it kept me from getting out of bed. One morning, I could not pull myself up and had to crawl. My MRI showed that disc L4–L5 was herniated. I started Non-Surgical Spinal Decompression, and by my fourth session, the pain was gone! At the end of my twelfth session, my flexibility was back to such a degree that I had not experienced for almost 12 years. One month previous to writing this letter my future looked bleak, with only invasive options to consider. I don't even want to think about what would have happened if I had not heard from a friend about Non-Surgical Spinal Decompression

— **Kent L.** —

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information.

## What Can I Expect?

### Getting Started



At your first visit, your doctor will conduct a physical exam and might recommend an X-ray or MRI to pinpoint the specific areas of damage and discomfort. Your doctor will then determine your course of therapy and whether you are a candidate for Non-Surgical Spinal Decompression (NSSD).

### The Pressure Is Off!

At the beginning of each session, you will be comfortably fitted with a pelvic harness designed to achieve optimal decompression of the lumbar spine. As a session of NSSD commences, you will notice a slow, gentle lengthening in your spine as your discs are gradually decompressed and relieved of pressure.



The treatment process is safe and relaxing. While some patients with extensively injured discs have reported mild discomfort during the first few treatment sessions, their discomfort subsides upon subsequent visits.

A patient safety switch, or quick-release clasps on the upper-body harness provide an extra safety feature, allowing you to stop at any point should you feel discomfort. Each treatment session lasts approximately 30–45 minutes.


\* Individual patient results may vary.

Copyright © 2005–2009 MediaWest. All Rights Reserved. [www.medwestpublications.com](http://www.medwestpublications.com)

### Typical Treatment Regimen

A typical NSSD treatment regimen consists of about 20–25 sessions over four to six weeks. Some conditions require fewer visits; some require more. Many patients report relief from their pain and other symptoms during the first few treatment sessions, and most experience dramatic pain relief after completion of their prescribed NSSD program.\*

As a session of Spinal Decompression progresses, the discs are relieved of pressure, creating a vacuum of negative pressure within the disc. Many experts believe this decompressed state aids in pulling nutrients, oxygen, and moisture back into the discs.



Patients get to lie back and relax while spinal pressures reach negative levels. Now the natural repair mechanism in the body has a chance to really do its work.

# What Is Causing My Back Pain?

While only a trained medical professional can accurately diagnose and recommend appropriate treatment, a basic understanding of common causes of back pain and how Non-Surgical Spinal Decompression works to alleviate them can help you in making a more informed decision concerning your treatment options.

## Understanding How the Spine Works


Your spine is composed of 24 bones called vertebrae. In between each vertebra is a fibrous disc (*annulus fibrosus*) filled with a jelly-like substance (*nucleus pulposus*), which provides flexibility and cushioning for the spine.

The vertebrae protect the spinal cord, which runs through a tube at the back of the spine called the spinal canal. In the lower portion of the back, spinal

nerves exit the spinal canal between the vertebrae and unite as they move down through the pelvis. Some of these spinal nerves join to become the sciatic nerves, which travel down through the buttocks, along the backs and sides of the thighs and calves, and into the feet.

With such a dense network of nerves traveling throughout the back, it's easy to see how great discomfort may be caused by a slight upset in the delicate architecture of the spine. Accidents and injury may damage discs and vertebrae, putting pressure on nerves. This results in tingling, numbness, muscle weakness, or even sharp, shooting pain.

Shown in the illustration below are some of the more common diagnoses for back pain that will be covered at length in the coming pages.



## Consider These Facts on Back Pain:

- On any given day, 6.5 million people are in bed because of back pain.
- Approximately 5.4 million Americans are disabled annually due to back pain.
- Back pain is the #2 reason for hospitalization.
- Up to 85% of the U.S. population will have back pain at some time in their life.
- After cold and flu, back pain is the number one cause of work absence.
- Spine surgery is the second leading surgical procedure in America with the total number in the U.S. approaching 500,000 per year.
- An estimated 93 million workers are lost each year due to back pain.

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.urmcmedwestpublications.com](http://www.urmcmedwestpublications.com) or call 877-968-8631 for ordering information.

## Lift with Your Legs, Not with Your Back!

Many bulging and herniated discs can be attributed to lifting objects incorrectly. When picking up an object from the floor, bend at the knees, keeping the spine as erect as possible. Rise using the strength of your legs, as their simple and sturdy architecture is designed for such exertions. Putting great amounts of pull or pressure on the spine when bent forward or backward can compromise its delicate alignment.

Of course, there is no right way to lift an object that is simply too heavy. Listen to your body, and if you feel any discomfort while doing heavy lifting, back off and save your back!



## Other Common Causes of Back Pain:



## "For three very long years

I had severe lower back pain. Doctors tried epidurals, to no avail, and said I would have to learn to live with it. I'd been using a walker to get around and could not stand more than five to ten minutes without excruciating pain. My activities were very limited. I was unable to sleep in a bed for three years. After reading an article on Non-Surgical Spinal Decompression Therapy, I decided to try it out.

After 12 visits, I am now sleeping in a bed, and my walker is put away. I'm able to go shopping for over an hour without any increased pain. I'm cooking meals and cleaning without having to sit down every five minutes because of the pain. My family cannot believe the changes. I thought I would have to live with this for the rest of my life. I am so thankful that I tried this and would encourage you to do the same!"

—Mardella S.—


# Bulging and Herniated Discs

Discs are located between each vertebra and provide flexibility and shock absorption for the spine. The thick, fibrous outer disc wall, known as the annulus fibrosus, surrounds a jelly-like center, called the nucleus pulposus. Discs undergo tremendous amounts of stress, which can sometimes lead to a bulging disc, which is when the disc wall becomes weakened enough to allow the disc to bulge out and press painfully on surrounding nerves. A herniated disc occurs when the pressure within a disc is so great that a tear occurs in the disc wall (annulus fibrosus), allowing a portion of the nucleus pulposus to protrude. The escaped nucleus pulposus may then impinge painfully on nerve roots, leading also to numbness, tingling, and/or muscle weakness associated with the condition of sciatic pain. The illustration below gives a bird's eye view of what exactly is going on when a disc bulges or herniates.



**The Jelly Donut Analogy**

Comparing the discs in your back to jelly donuts gives you a good idea of what happens when a disc herniates. Picturing in your mind a jelly donut plump with strawberry filling. Have you ever set a box of donuts on your office chair, only to absentmindedly plip down on them a few moments later? What happens? The jelly oozes out under the pressure. When a disc ruptures or herniates, the same thing happens, except, unlike the jelly donut, a herniated disc can be repaired! That's where Non-Surgical Spinal Decompression comes in.



# Sciatica, Spinal Stenosis


## What Is Sciatica?

Sciatica is the sensation of pain, tingling, or numbness in the buttocks and/or legs produced by an irritation of the sciatic nerve. Multiple nerve roots extend from the spinal cord between the vertebrae and join to form the sciatic nerve, which branches as it descends into the buttocks and down each leg to the ankles and feet. The primary causes of sciatica are herniated, bulging, or degenerated discs, which induce pressure on the spinal nerve roots.

Other causes include small, bony growths on the spine (bone spurs) or compression of the nerves through injury.


In rare cases, the sciatic nerve may be irritated by conditions such as tumors, pregnancy, or piriformis syndrome.

- L1** ■
- L2** ■
- L3** ■
- L4** ■
- L5** ■
- S1** ■



## Where Does It Hurt?

In the illustration below, the letters L1–L5 refer to the lumbar (lower) portion of the spine and S1 refers to the first segment of the sacral vertebrae. Over 90% of herniated discs occur in the lowest two levels of the lumbar spine, between L4–L5 and L5–S1. The colors reflect the regions of the lower body potentially affected by compression of each given lumbar and sacral nerve root. Do you experience pain, tingling, or numbness in any of these regions?



## What Is Spinal Stenosis?

Stenosis is a narrowing of the spinal canal due to the encroachment of disc material or bony growths that squeeze and irritate the spinal cord and/or extending nerve roots. This can lead to pain, numbness, tingling, or weakness in the legs, feet, or buttocks. The benefits that many stenosis patients derive from Non-Surgical Spinal Decompression may be due to its positive repositioning and rejuvenating effect on the herniated and degenerative discs that often accompany stenosis. Spinal stenosis, which may be found in conjunction with any of the above conditions, is commonly a contributing cause for sciatic symptoms.



# Degenerative Disc Disease

## What Is Degenerative Disc Disease?

Degenerative disc disease is not technically a disease, but rather a state of disc dehydration and deterioration due to a combination of cumulative trauma, poor dietary and exercise habits, and aging. As discs degenerate they become more prone to failure from physical stress, which may tear disc fibers and result in more complications, such as osteoarthritis, disc bulging, disc herniation, and stenosis.

Many spine experts conclude that the vacuum of negative pressure created within the disc by Non-Surgical Spinal Decompression helps the disc to attract moisture from surrounding tissue, rehydrating and revitalizing thinking and torn degenerated discs.\*

## If You Don't Use It, You'll Lose It

Traumatic spinal injuries may cause patients to avoid their normal daily activities. Without proper treatment, pain will progressively worsen, resulting in decreased physical activity and gradual weakening of the supportive spinal muscles, leading to accelerated disc degeneration.

## Going, Going... and Finally Gone

### Phase I: Dystunctional

Phase I of degenerative disc disease is categorized by tears around the outer surface of the annulus. Further damage to the disc and surrounding tissue is exacerbated by the less effective disc.



©MediaWest

### Phase II: Unstable

In Phase II, the joint progressively loses strength. Disc changes include further tearing along the horizontal side of the disc, greater loss of disc height, and cartilage degeneration.



©MediaWest

### Phase III: Stabilization

Further loss of disc height, disc space narrowing, moderate to severe endplate tearing along the horizontal side of the disc, greater loss of disc height, and surrounding vertebrae to fuse together.



©MediaWest

\* Individual patient results may vary.

"I'm eighty-nine years old and I have had low back pain off and on for years. For the past eight months I had terrible back pain every single day. Amazingly, after only six treatments of Non-Surgical Spinal Decompression, I no longer needed pain medications or the use of a cane to get around my home or garden."

— Margaret S. —


"After two years of dealing with the pain, and, at times, hardly being able to walk with even a cane, I am so grateful to say that I am finally pain free. Non-Surgical Spinal Decompression was truly an answer to my prayers. Thank you so much for fixing my back."

— Earlene O. —

# Facet Syndrome

## What Is Facet Syndrome?

Facets are the bony, wing-like protrusions extending from the back of the vertebrae that align with facets on the vertebrae above and below, and give the spine a more diverse range of motion. Facets function as guides for the spine and are not designed for bearing weight like vertebrae and discs. Joining adjacent facets are small ligaments called facet joints. Facet syndrome involves inflammation of the facet joints. It is one of the lesser-known but surprisingly common causes of back pain.



©MediaWest

## Bearing the Brunt of Unhealthy Discs

When a person is standing upright, the facet joints bear about 16% of the normal compressive forces of the spine. With disc height loss of 1-3 mm, the compressive load can be five times the normal amount on the facet joints.<sup>1</sup> Bearing the brunt of all that weight can lead to tearing or degeneration of the ligaments, as well as inflammation of surrounding tissues. Adhesions over the joint surface can form over time, leading to loss of mobility and breakdown of facet cartilage. An indicative symptom is a deep ache in the lower back that may extend to the buttocks, hip, and even below the knee. Facet syndrome is often associated with degenerative disc disease and soft tissue damage in the lumbar spine.

## Contributing Factors to Spinal Injury and Disease



**Age:** As we grow older, discs may dry and crack, losing flexibility and the ability to cushion the vertebrae. Good diet, plenty of exercise, and water intake can help slow the process.



**Exercise:** Exercise keeps the muscles surrounding the spine strong, decreasing injury susceptibility. Exercise also helps to maintain healthy blood flow to discs and surrounding tissue.



**Diet:** A healthy diet with appropriate supplementation will help ensure that your bones and tissues are receiving proper nutrients.



**Activities:** Maintaining an awareness of what your body can handle. Falls or reckless exertion on the body, such as lifting heavy objects, can lead to severe spinal injuries. Be smart!

1. M.A. Adams, M.C. Hutton. The effect of posture on the load of the supporting joints in standing. *International Orthopaedics*. 1980;6:288-92.  
2. Charles H. Adams, M.D., M.D., M.C. Disc space narrowing and the lumbar facet joints. *J Bone Joint Surg*. 1984; 66:704-10.

## Stories of Recovery: Patients Talk



"I had surgery for the repair of a ruptured disc in 1997. Over a decade later, I once again found myself in agonizing pain. The doctor I was seeing told me that surgery was my only option. After seeing the amazing results that my husband had using Non-Surgical Spinal Decompression—he suffered from low back pain for 46 years and was now pain free—I was hoping the procedure would also help me. I was in so much pain when I came to the doctor's office that I needed the help of two people just to walk up the ramp. I was hoping and praying for a miracle. After only two weeks of treatment, my pain subsided dramatically and, after finishing the program, I am now pain free. It's absolutely amazing. I cannot believe how great I feel! Can you do anything now without pain. This was the answer to my prayers." — **Lilian C**



"I was hospitalized for 30 days, bed-ridden with excruciating low back pain, due to a ruptured disc. I began researching on the Internet for an alternative to surgery, and found out about Non-Surgical Spinal Decompression. I decided I really didn't have anything to lose. After completing the full treatment regimen, I had found the answer to my chronic low back pain. The post-treatment MRI showed that my ruptured disc had 'unruptured' and repaired itself! Thank you for giving me back my life!" — **Richard H.**



"I arrived at my first appointment in terrible pain. Before Non-Surgical Spinal Decompression, I had trouble sleeping, getting dressed, taking a shower, and even performing simple tasks at work. Two months later I am 99% better and can do the things I used to." — **Addie K**



"I suffered for years with low back pain from herniated discs, stenosis, arthritis, and bone spurs. After having tried pain management, epidural injections, physical therapy, and meds, I was very skeptical about Non-Surgical Spinal Decompression. Boy, was I surprised! I am totally pain free, very active, and I tell everyone about this wonderful therapy!" — **Edward D.**

"I injured my back while kayaking. An MRI showed I had stenosis and a herniated disc, which caused pain and sciatica in my left hip and leg. I tried massage, chiropractic, acupuncture, exercises, and rest. I feared things would never get better and that I'd never enjoy an active life again. Then friends suggested Non-Surgical Spinal Decompression. In the waiting room, other clients shared their success stories, and I had hope for the first time in a year. My recovery was like many others. After 15 treatments, my pain was gone. I had a few more treatments and was soon back to being able to work, garden, hike, ride my bike, kayak, and otherwise enjoy my life to the fullest without pain. Now, 18 months later, the pain has not returned. I can't say enough about how grateful I am." — **Barbara D.**

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Please visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information. Copyright © 2005–2009 MediaWest. All Rights Reserved.

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Please visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information.

## Doctors Talk



"As an interventional pain physician, I feel that Non-Surgical Spinal Decompression has filled a void in our continuum of treatment for those patients suffering from back and neck pain associated with disc pathology. The majority of the other procedures I perform, such as injections and nerve blocks, treat the pain due to tissue and nerve irritation, which only reduce the patient's pain. My enthusiasm for Non-Surgical Spinal Decompression stems from the fact that it is the only therapy that scientifically attempts to heal and rejuvenate the discs. My personal experience with this noninvasive therapy is that qualified patients will show improvement within the first few treatments." — **Dr. Juan J. Gargallo, MD** Physician



"Most patients we see have very difficult, debilitating low back conditions that could not be helped by conventional pain management. After 20,000 Non-Surgical Spinal Decompression treatments over the last six years we've seen some pretty amazing things. The most satisfying is to see most of our patients stay out of pain years later. Our first two patients that we treated back in 2003 are virtually still pain free to this day!" — **Dr. Kenneth Zaminillo, DC** Chiropractic Physician



"Non-Surgical Spinal Decompression Therapy has been an extremely valuable tool in caring for patients that many times have already had physical therapy and are now being told that surgery is a consideration. Of this population, I'm easily fixing the majority of even the most difficult chronic patients." — **Dr. Francisco Ward, DO, DABPMR** Osteopathic Physician



"I've had the pleasure of helping hundreds of patients with chronic back problems who thought there was no hope. Finally, a treatment has surfaced that directly treats the damaged tissue that is causing the pain. Many of my patients have previously undergone unsuccessful treatments with physical therapy, painful injections, and/or surgery. I would never practice again without spinal decompression." — **Dr. Christopher R. Houtakker, DC, CCST** Chiropractic Physician



"Many new patients come in wary and skeptical because of past negative experiences with some member of the health community. Non-Surgical Spinal Decompression Therapy provides instant enthusiasm and immediate receptivity. You can see the light of hope in their eyes. My Decompression Therapy unit is the cherry on top of all the various tools used in my practice!" — **Dr. Pierre Parent, DC** Chiropractic Physician

# How Spinal Decompression Works


High intradiscal pressures cause discs to bulge out and press painfully on nerve roots. They also make for a compressed, anaerobic environment unsuitable for healing.

Non-Surgical Spinal Decompression (NSSD) produces negative pressures within the disc, which experts believe create a vacuum effect that draws in nutrients and fluids to promote the repair of injured discs and surrounding tissues. This vacuum has also been shown to aid in the retraction of escaped cushioning gel from herniated discs.

## When Negative Is a Positive

Much like gauging the air pressure in a car tire, scientists can use sensors to measure the various pressures put on spinal discs while lifting, standing, sitting, lying down<sup>1</sup>, undergoing traction<sup>2</sup>, and undergoing NSSD. Like other pressures found in the body such as blood pressure, intradiscal pressure is measured in millimeters of mercury (mm Hg).

While traction, physical therapy, and manipulation may reduce disc pressures to as low as 40 mm Hg, only NSSD has been shown to achieve negative pressures within the spine. It has been clinically proven that NSSD creates negative pressures as low as -160 mm Hg<sup>3</sup> within the injured disc during the treatment session!



## Decompression Goes Beyond Normal Traction

With traction, weights are steadily added to the end of the traction bed, which in turn, adds tension to a harness secured around the patient's pelvis, lengthening the spine. The intention is to relieve pressure, but the linear force can instead produce spasming, which may lead to greater injury.

A 1997 study<sup>4</sup> compared a traditional mechanical linear traction device to a Non-Surgical Spinal Decompression device. The traction device yielded no positive results with herniated discs, while the NSSD device yielded "good" to "excellent" results. In the vast majority of patients treated, like traction, NSSD also lengthens and exerts tension upon the spine, but the approach is far different, and produces vastly superior results.

## The Logarithmic Difference

Normally, pulls exerted on the spine trigger sensory receptors in the back to tighten the muscles surrounding the vertebrae and discs in an effort to protect them from injury—a mechanism in the body known as the proprioceptor response. NSSD bypasses this response by gently pulling on the spine and relaxing the back over an extended period of time, allowing the spine to be repositioned without tension, and without setting off the "lock down" proprioceptor response.


1. Aftabzadeh, MD, PhD, The Load on Lumbar Discs at Different Postures of the Body, *Chiropractic*, 55, 107-122, 1986.  
 2. Goh, Andrew, MD, PhD, Aftabzadeh, MD, PhD, University of the Research Center for Traction, *Spine*, 19, 1111-1116, 1994.  
 3. Goh, Andrew, MD, PhD, Aftabzadeh, MD, PhD, University of the Research Center for Traction, *Spine*, 19, 1111-1116, 1994.  
 4. Cramer, Stanley, MD, PhD, Van Bortom, Ph, MA, *Chiropractic*, Education, and Rehabilitation of the Lumbar Spine & Cervical Traction Treatment for Lumbosacral Pain, *American Journal of Physical Medicine*, Vol. 7, No. 2, April 1977, Last page 21 for research details.

# The Key to Spinal Decompression

## Formula for Relief

Each automated session of Non-Surgical Spinal Decompression (NSSD) cycles the patient through a series of gentle pulls, holds, and releases. Super-smooth transitions between each phase of NSSD can make for an experience so relaxing that patients often fall asleep!

As each session commences, tension slowly mounts, lengthening the spine. Up to one-half of the patient's body weight, plus as much as 25 pounds of tension, can be exerted directly on the injured discs—all without triggering the "guarding" proprioceptor response. This is where spinal pressures drop and decompression actually occurs. The differing amounts of tension administered throughout the session, when graphed on a chart, produce a logarithmic curve similar to the one shown above. After a holding period, tension is slowly decreased and the spine is gradually retracted. This cycle is repeated several times throughout a treatment session.



## Before Non-Surgical Spinal Decompression



## During Non-Surgical Spinal Decompression

NSSD helps reposition extruded disc material back into place. Pressure on spinal nerves is reduced.



1. Individual patient results may vary.  
 2. Mitchell Schramke, MD, PhD, Ed. Stanley, MD, Udo Schramke, MD, *Theme Atlas of Anatomy*, Germany, Georg Thieme Verlag, 2006.



# Is Spinal Decompression Right for Me?

## You Are a Candidate for NSSD If:

- You have chronic or severe back pain caused by bulging or herniated discs, degenerative disc disease, sciatica and/or facet syndrome.
- You have failed back surgery syndrome.
- You have been told to consider surgery.

## You Are *Not* a Candidate for NSSD If:


- You have any retained surgical hardware (titanium rods or fusion cages).
- You have been diagnosed with a clinically unstable back (spondylolisthesis).
- You have rare conditions such as certain spinal infections, or pelvic or abdominal cancer.
- You have sustained a recent vertebral fracture.
- You suffer from severe osteoporosis (DEXA T-score of -2.5 or lower).
- You have ankylosing spondylitis.
- You have an abdominal aortic aneurysm.
- You are pregnant.

## When Can I Expect Results?

Many patients report a reduction in pain after their first few NSSD treatment sessions and happily feel they have had all the treatments they need. This is a wonderful sign that NSSD is working, but the scientific reality is that the healing process is not finished! Quitting the treatment regimen early usually results in a relapse of symptoms, as well as wasted time and money.

On the other hand, it may take several sessions before patients experience a noticeable remission of symptoms.

• Individual patient results may vary.



Why is this? Bulging and herniated discs may need several sessions to fully reposition themselves depending on your physiology. You can be sure that your spine is responding to NSSD's vacuum effect, but results are not always immediate. Using time-tested protocols, NSSD can be tremendously effective at eliminating back pain!

## NSSD Works, Given the Chance

Some patients report an immediate pain reduction after their first treatment! With herniated and degenerated discs, the escaped nucleus pulposus may be partially or completely retracted back into the disc after just the first session (though it normally takes longer), relieving a great deal of pain.\*

However, pain reduction does not indicate full recovery, and a full recovery is essential to preventing relapse. This may mean completing as many as 20 to 25 treatment sessions. Older or more severe injuries may require even more sessions. This may seem like a lot of treatments, but NSSD is working with your back as it heals, and that is a process that cannot be rushed. Why is this?

## Follow Through for Best Results

As with all wounds, tears in the disc wall need time to heal. The absence of pain does not mean the tears have had time to seal up, which is crucial to preventing the newly retracted nucleus pulposus from escaping and putting pressure back on the nerves.

# Completing Your Course of Therapy

Completing your course of Non-Surgical Spinal Decompression (NSSD), as determined by your doctor, requires commitment to feeling good again. The time needed to complete a treatment course varies according to the severity of your condition. Each session prescribed is needed to maintain a fully hydrated and oxygen-rich environment for the damaged disc. The same principles apply for those with degenerative disc disease.


## Visible Progress!

After as little as nine treatments, the injured disc wall may have almost completely healed. However, a small fissure remains. This fissure still requires more sessions of NSSD in order to properly mend itself. Continuing on with the NSSD protocol will allow for an enriched disc environment to speed the body's natural healing mechanism. The last remaining NSSD sessions will enable the disc to heal completely.

## "I have tried many remedies

In an effort to find relief from back pain that plagued me for two years, the pain from my sciatic nerve would be so bad at times I was in tears. Often times I couldn't find a comfortable position even while lying down. When I thought I was going to have to learn to live with the pain, I discovered Non-Surgical Spinal Decompression Therapy and, at last, relief! I've had 16 treatments and I'm finally feeling improvement! Before the treatments, I could only stand for ten minutes without pain. Now I can stand without pain, and everyday activities are easier!"

—Marilyn B.—



**1** Before NSSD: The annulus has torn under pressure, and the escaped nucleus pulposus presses painfully on the sciatic nerve, symptoms include numbness, tingling, and pain.

**3** It's Working! The nucleus pulposus may be completely repositioned within three to five treatments. Blood, oxygen, and nutrients flow into the disc, aiding in the healing of the torn annulus.



**2** First Signs of Relief: In as little as one or two treatment sessions, the nucleus pulposus may be retracted into the disc, taking pressure off nerves, thus dramatically reducing pain.



**4** Finished! With the newly repositioned nucleus pulposus, patients can return to a normal, healthy, and active lifestyle!

Note: These artistic renderings offer a simplified explanation of complex physiological processes.

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information. Copyright © 2005-2009 MediaWest. All Rights Reserved.

## Before Surgery, Consider This!

While back surgery may sometimes be the only solution, many medical professionals and surgeons themselves feel strongly that every noninvasive option should be explored before turning to surgery.

Although advances in surgery have made many procedures less invasive and more effective, surgery does come with inherent risks. With high costs, lengthy recovery time, and possible infection, one should consider all options before making a decision about surgery.



### Failed Back Surgery Syndrome

Failed Back Surgery Syndrome is a real term used when a patient continues to suffer from pain and loss of mobility long after surgery. According to the American Academy of Orthopedic Surgeons, there are approximately 200,000 laminectomies performed every year with an estimated 20-30% of these operations reported to be unsuccessful. A laminectomy removes bone and thickened tissue that is narrowing the spinal canal and squeezing the spinal cord and nerve roots.

### Great News for Post-Surgical Patients!

If you still suffer from pain after surgery, we have great news for you! Post-surgical patients have experienced fantastic therapeutic results from Non-Surgical Spinal Decompression (NSSD).<sup>1</sup> While those with surgical hardware<sup>2</sup>, such as brackets, plates, and screws, are not candidates for NSSD, many post-operative patients are. If you have had surgery with unsatisfactory results or have suffered a relapse, consult your doctor about the possible benefits of NSSD.

### "There is a place for surgery



but it is definitely not at the front line. Non-Surgical Spinal Decompression Therapy should be a front line treatment for chronic back pain patients suffering from bulging, herniated, or degenerative disc disease and facet syndrome.

As a surgeon, I only want to do surgery when I absolutely have to. Spinal Decompression Therapy gives my patients a more conservative treatment option that can eliminate the need for surgery altogether, and that's a very good thing.

Non-Surgical Spinal Decompression Therapy also gives me a treatment option for those patients who have had surgery and have a relapse or experience a return of pain following surgery. It allows me to offer patients a key, non-surgical weapon in the war on back pain.

— **Dr. Bernard Zaliger, DO** —  
**FACOS, FAOAO, FICS**

Osteopathic Physician and Orthopedic Surgeon  
Founding Dean and Provost of Tuoro University  
College of Osteopathic Medicine, Vallejo, CA

1. Individual patient results may vary.

2. Although surgical hardware is a contraindication for Non-Surgical Spinal Decompression, post-surgical patients with low back hardware may qualify for cervical treatment, and patients that have had surgery with hardware in the cervical area may qualify for lumbar treatment. This is to be determined on an individual basis.

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MediaWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.medwestpublications.com](http://www.medwestpublications.com) or call 877-968-8631 for ordering information.

## Dramatic Results Without Surgery

### Case Study: Severe Herniated Disc

Two years ago, Terence M. ruptured a disc while lifting a box from the trunk of his car. As a result, he lost the use of his right leg and suffered relentless, excruciating pain. Desperate for relief, he went to see Harvey Kleinberg, DO, an Osteopathic Physician and Physical Medicine and Rehabilitation Specialist for over 35 years. Dr. Kleinberg used Non-Surgical Spinal Decompression Therapy to alleviate Terence's pain and restore his lost function.

"When patients have injuries, you want to make them feel better as quickly as possible, and that is what Spinal Decompression Therapy is capable of doing for many patients," Dr. Kleinberg said.



Dr. Kleinberg prescribed a treatment plan of three 30-minute sessions a week, which eventually tapered off as the pain began to subside. Patients usually undergo about 20 to 25 treatments, and easy maintenance can keep people from ever having a repeat injury again.

"My pain was excruciatingly horrible," Terence said. "I thought I'd never be able to be up and about again. My pain was gone in two weeks; I was feeling so good, I couldn't wait to go back to work."

### "I Was a Pain Doctor In Pain!"

Dr. Kleinberg knows first-hand how unbearable back pain can be—he suffered from herniated discs three years ago. "I had undergone three laminectomies, but continued to suffer from severe sciatica one year after the third surgery. Due to the severity of my problem, it took 29 treatments for a total elimination of pain. It's been three years now, and I'm still pain-free!" Because Spinal Decompression Therapy alleviated his back pain, he is determined to inform people about this non-surgical treatment for herniated or degenerative discs.

"No one should have to suffer from pain for the rest of their lives if there is a treatment outside of surgery that can offer them relief with no risk involved," Dr. Kleinberg said.



### Pre- and Post-MRIs: Spinal Decompression in Action!

Shown in this box are pre- and post-MRIs of a non-surgical Spinal Decompression Therapy patient who had a herniated disc much like that of Dr. Kleinberg's patient, Terence. The red arrows point to the injured disc, before and after treatment. Before treatment, the disc protrudes into the spinal canal, impinging painfully on sensitive nerves. After treatment, the once-extruded disc material has been literally "sucked" back into place! The result was a complete return to normal functioning.\*

\* Individual patient results may vary.

# CAUTION

**Pain Medications—Special Advisory Warning: If you manage your pain medication intake. In the U.S., 60 million adults take Over-The-Counter OTC pain relievers every day or for several days per week.<sup>1</sup> Ongoing use can lead to serious health risks or even death. OTC pain drugs should be taken at the lowest effective dose for no more than 10 days.<sup>2</sup>**

## Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)

This class of drug is the most commonly used medication in the world. OTC NSAIDs include Ibuprofen (Advil<sup>®</sup> and Motrin<sup>®</sup>), naproxen (Aleve<sup>®</sup>), and aspirin. NSAIDs are used primarily to treat inflammation, pain, and fever, by blocking prostaglandin production. However, prostaglandins are needed to create mucosal protection for the lining of the stomach, and, if blocked, serious damage can occur. NSAIDs also inhibit platelets that help with blood coagulation and homeostasis, and can interfere with kidney function. There is no risk-free NSAID dose (includes low-dose aspirin).

## BLACK BOX WARNINGS

The FDA's most serious warning must now appear on all NSAID packaging.<sup>3</sup> This means the first time ever that an OTC drug has been required to have a Black Box Warning.<sup>4</sup> High doses of OTC brands can be just as dangerous as prescription formulations.

■ NSAIDs (except aspirin) may cause an increased risk of heart attack, blood clots, and stroke, which can be fatal. Risks may increase with duration of use. Cardiovascular conditions such as high blood pressure may also increase the risk.<sup>5</sup>

■ NSAIDs increase the risk of serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal and most often occur without warning symptoms.<sup>6</sup>

**NSAID-Induced PUEs**, known together as Perforations (erosions of the stomach and small intestinal lining) ulcers, and GI Bleeds, cause over 100,000 hospitalizations annually.<sup>7</sup> Chronic NSAID use (90+ days) showed serious damage to the small intestine in 70% of the subjects.<sup>8</sup> Patients over 60 with perforations have a 30% chance of dying.<sup>9</sup>

Compare to non-NSAIDs, NSAID users' risk for PUEs increases by:  
2.5 times when taking low-dose aspirin.<sup>10</sup>  
3 times when taking OTC naproxen or rofecoxib.<sup>11</sup>

10 Times when taken with alcohol.<sup>12</sup>  
12 Times when taken with aspirin.<sup>13</sup>  
12 Times when taken with aspirin and alcohol.<sup>14</sup>

13 times when patients report liver disease or GI Bleed.<sup>15</sup>  
13 times when used with anticoagulants (blood thinners).<sup>16</sup>

15 times when used in the cardiovascular.<sup>17</sup>  
25-30 times when given NSAIDs are used together.<sup>18</sup>

**Acetaminophen (APAP)** Tylenol<sup>®</sup>, Daitil<sup>®</sup>, and others do not have anti-inflammatory effects like NSAIDs, but are commonly taken for chronic pain. Used in over 600 medications.<sup>19</sup>

➤ Annually, acetaminophen (APAP) toxicity kills nearly 500 people and causes 55,000 ER visits, 2,500 hospitalizations, and 100,000+ calls to Poison Control Centers.<sup>20</sup>

➤ Overdose of acetaminophen is the leading cause of **Acute Liver Failure (ALF)**. ALF may feel like flu symptoms over several days.<sup>21</sup> Coma and death can rapidly occur in one-third of ALF cases.<sup>22</sup>

➤ ALF can occur using the maximum 4 grams (gm) per day dose for five or more consecutive days.<sup>23</sup> 10% of ALF victims used 2-4 daily gm.<sup>24</sup>

➤ APAP causes half of all ALF. Of these cases, 38% had combined two or more APAP-containing preparations.<sup>25</sup>

➤ Alcohol used with more than 2 gm of APAP can cause ALF.<sup>26</sup>

## Bigger Doses Don't Mean Better Relief

The analgesic ceiling effect of a drug refers to the dose beyond which there is no additional pain relief. Taking higher than the recommended dose does not yield additional pain relief, but may increase side effects. The ceiling per dose for the following meds are: Ibuprofen, 400 mg; Acetaminophen, 1000 mg; Naproxen, 500 mg; Pure Opioids, no ceiling; Combination Opioids with APAP (less than 4 gm/24 hours to avoid ALF).<sup>27</sup>

**Opioids** These powerful prescription narcotics are extremely addictive and may cause permanent physical changes in the brain. Commonly prescribed opioids are oxycodone (OxyContin<sup>®</sup>), hydrocodone (well-known brands Vicodin<sup>®</sup> and Lortab<sup>®</sup> contain acetaminophen), and methadone.

➤ The #1 selling U.S. drug is hydrocodone. With 135 million prescriptions, the U.S. uses 99% of the world's hydrocodone.<sup>28</sup>

➤ Hydrocodone caused 62% of accidental APAP-induced ALFs.<sup>29</sup>

➤ Prescription opioid-related deaths exceed the total deaths involving heroin and cocaine.<sup>30</sup>

➤ There were 22,400 accidental overdose deaths with opiate prescriptions vs. 17,000 homicides in 2005 in the U.S.<sup>31</sup>

➤ More people in the 45-54 age group die of unintentional opioid drug overdose than from motor vehicle accidents.<sup>32</sup>

# Clinical Studies & Publications\*

This document is a sample brochure. It is copyright protected by the Library of Congress. No portion of this booklet may be shared or used in print, electronic, or web-based media without the express permission of MedWebWest. Full-color, hard copy versions may be purchased for patient educational purposes. Visit [www.medwebwestpublications.com](http://www.medwebwestpublications.com) or call 877-956-8531 for ordering information. Copyright © 2005-2009 MedWebWest. All Rights Reserved.

## American Journal of Pain Management: Long-term Effect Analysis of IDO THERAPY<sup>®</sup> in Low Back Pain: A Retrospective Clinical Pilot Study, July 2005, Vol. 15, No. 3, C. Norman Shealy, MD, PhD, Norman Kotzick, MD, Merrill M. Weisman, MD.

**Outcome:** Of 24 study participants, each reported consistent pain relief and continued improvement of symptoms one year later. Improvement in pain continued after the treatment sessions were completed.

## Practical Pain Management: Technology Review: IDO THERAPY<sup>®</sup>, April 2005, Vol. 5, Issue 3, C. Norman Shealy, MD, PhD.

**Outcome:** The treatment leads to satisfactory pain relief and improved quality of life in up to 88% of patients—many of whom have failed other "conventional" approaches. Based on the author's review of recent study results, IDO THERAPY<sup>®</sup> appears to be the current optimal recommendation for most lumbar pain syndromes.

## Journal of Neuroimaging: MRI Evidence of Nonsurgical, Mechanical Reduction, Hydration and Repair of the Herniated Lumbar Disc, April 1998, Vol. 6, No. 2, Edward L. Eberman, MD.

**Outcome:** All but 3 of 20 patients reported significant pain relief and complete relief of weakness and immobility, when present. This study also shows a correlation between the improvement on the MRI and the reported improvement in pain.

## Journal of Neurological Research: Vertebral Axial Decompression for Pain Assoc. with Herniated or Degenerated or Facet Syndrome: An Outcome Study, April 1998, Vol. 20, No. 3, E. Gose, PhD, W. Noguyszewski, MD, R. Noguyszewski, MD.

**Outcome:** Pain, activity and mobility scores greatly improved for 71% of the 778 patients studied. The authors consider VAXD<sup>®</sup> to be a primary modality for low back pain due to lumbar herniations, degenerative disc disease, and facet arthropathy. The authors concluded that post-surgical patients with persistent pain or "Tailed Back Syndrome" should try VAXD before further surgery.

## American Journal of Pain Management: Decompression, Reduction, and Stabilization of the Lumbar Spine: A Cost Effective Treatment for Lumbosacral Pain, April 1997, Vol. 7, No. 2, C. Norman Shealy, MD, PhD, Vera Bogomyey, RN, MM.

**Outcome:** The authors compared the pain-relieving results of traditional mechanical traction (14 patients) with a decompression device (25 patients). The decompression system gave "good" to "excellent" relief in 85% of patients with ruptured discs and 75% of those with facet arthrosis. The traction yielded no "good" to "excellent" results in patients with facet arthrosis.

## Journal of Neurosurgery: Effects of Vertebral Axial Decompression on Intradiscal Pressure, September 1994, Vol. 81, No. 3, Gustavo Ramos, MD, William Martin, MD.

**Outcome:** VAXD<sup>®</sup> creates a negative intradiscal pressure force up to -160 mm Hg.

\*These studies were performed on different Non-Surgical spinal decompression devices using various treatment protocols on different patient populations with different study criteria. They do not necessarily represent whether results will be achieved with particular spinal decompression devices.

\*\*IDO THERAPY<sup>®</sup> is a registered trademark of North American Medical Corporation (NAMI) to be used only with certain certified NAMI devices by professional staff specifically licensed by NAMI.

†Note: The key clinical research documenting the negative pressure force created within "in vivo" (living) intervertebral discs during Non-Surgical Spinal Decompression (NSD) was performed using a VAXD<sup>®</sup> system. VAXD<sup>®</sup> is a registered trademark of VAXD Medical Technology, Inc. All NSD devices currently registered with the FDA have received this VAXD<sup>®</sup> clearance by claiming their device is substantially similar to produce traction devices.



Pre-Treatment MRI Post-Treatment MRI  
Property of PPI Communications  
Used with permission: PPI Communications, Inc.

1. C.A. Wilson et al. *Contemporary Management of Back Pain*, 2nd Edition, Lippincott Williams & Wilkins, 1994, p. 1002-1008.

2. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

3. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

4. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

5. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

6. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

7. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

8. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

9. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

10. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

11. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

12. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

13. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

14. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

15. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

16. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

17. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

18. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

19. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

20. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

21. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

22. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

23. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

24. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

25. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

26. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

27. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

28. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

29. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

30. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

31. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

32. <http://www.fda.gov/cder/rdmt/OTC/OTCNSAID.htm>

## More Patients Talk



"I suffered from low back pain with numbness and tingling in my left foot and leg after a lifting injury. Medical doctors told me they could try a nerve block or I could have surgery, and that was it. I definitely didn't want to have surgery, so I looked for an alternative. After seeing a commercial for Non-Surgical Spinal Decompression, I called to make an appointment. I was very skeptical at first, but after three months of treatment, my back pain, tingling, and numbness that I've had to live with for over 10 years is gone." — **Dan M.**



"I had to use a walker to get around. An MRI showed I had spinal stenosis, and the herniated discs in my lower back were pinching the sciatic nerve. I was having excruciating pain day and night, couldn't sleep, and no amount of medication could stop the pain. I had read about Non-Surgical Spinal Decompression Therapy. After a few appointments, I was doing much better and able to walk without the walker or crutches. I only wish I had done this right away!" — **Joan G.**



"My life has completely turned around. I hurt myself at work and I had undergone two back surgeries but was still in constant pain. We were struggling to make ends meet and had to go to church for financial help. We were a clergy case. We had to constantly tell our kids no to activities they wanted to do. Now, 18 months after undergoing Non-Surgical Spinal Decompression, I am still completely pain free. I can now provide for my wife and kids, and we can afford to have our kids participate in the same activities as their peers." — **David H.**



"I had been feeling miserable since 1989 with back pain and pain down my left leg. I tried many treatments before, including steroid shots, acupuncture, shoe inserts—nothing worked. Finally, I found something that did work! Non-Surgical Spinal Decompression has been great. I'm now almost completely pain-free and ready to go dancing!" — **Mary L.**



"I used to have constant low back pain ranging from mild to severe depending on what I did. I had sharp, shooting pain down my leg and tingling into my left foot. After only a few treatments with Non-Surgical Spinal Decompression, the leg and foot symptoms were gone and the back pain lessened. I'm now about 80% better and can do things I haven't been able to do for years without pain or with very little pain. I am very thankful for my doctor and this therapy!" — **Virginia J.**

## More Doctors Talk



"As occupational medicine specialists, we see many people who suffer from chronic back and neck pain. Like many other physicians, we have been frustrated by those who do not respond to traditional approaches. With Non-Surgical Spinal Decompression, we now have a non-invasive therapy that works to correct the underlying causes of the pain, providing relief in the vast majority of people. This treatment can often help even the most difficult cases, including post-surgical patients, resume active lives. With tens of thousands of patients successfully treated without surgery, you've got to wonder about those who call this technology experimental! You can't have a state-of-the-art back pain program without this technology!"

**Dr. Ronald Klein, MD** (Physician) & **Dr. E. Robert Womack II, DO** (Occupational Physician)



"I have been treating patients for over five years with Non-Surgical Spinal Decompression with incredible results. I have never seen another procedure that can reproduce these kinds of results. I have great success in patients with disc problems whom I would otherwise have referred out for invasive procedures. In my opinion, Non-Surgical Spinal Decompression will become the standard of care for low back pain sufferers in years to come."

**Dr. Roger Roughley, D.C.** (Chiropractic Physician)



"We have treated thousands of patients who have experienced long-term, pain-free healing because of decompression therapy. With Non-Surgical Spinal Decompression we offer real modification of the patient's disc disease processes and rehabilitation of the neuropathic and mechanical symptoms, rather than just offering palliative care. I use it for myself, and it is worth its weight in gold!"

**Dr. Phil Fisher, D.O., PhD** (Chiropractic Physician)



"I've always considered myself to be in the 'results business.' As a doctor of chiropractic who specializes in disc-related problems, patients seek our care with a single purpose in mind—to get better. After seeing results on real patients, whom we have determined to be good candidates for Non-Surgical Spinal Decompression, I don't simply believe this treatment is effective—I know it is. I used to recommend this treatment to patients based on the research. Now I recommend it because of the results."

**Dr. Michael Rozenthal, D.C.** (Chiropractic Physician)



"Neurologists like myself have long known that we should do everything possible to help our patients avoid back surgery. Now with Non-Surgical Spinal Decompression, we finally have a very effective way to treat back pain without surgery. The vast majority of even our worst cases experience significant, long-lasting relief if they complete the entire treatment regimen."

**Dr. Orlando Maldonado, MD** (Physician)